

19



Europäisches Patentamt
European Patent Office
Office européen des brevets



11 Publication number:

0 469 477 A1

12

EUROPEAN PATENT APPLICATION

21 Application number: **91112577.1**

22 Date of filing: **26.07.91** ,

51 Int. Cl.⁵: **A61K 31/55, A61K 31/44,**
/(A61K31/55,31:425),
(A61K31/44,31:425)

30 Priority: **02.08.90 US 561743**

43 Date of publication of application:
05.02.92 Bulletin 92/06

84 Designated Contracting States:
AT BE CH DE DK FR GB IT LI LU NL SE

71 Applicant: **F. HOFFMANN-LA ROCHE AG**
Postfach 3255
CH-4002 Basel(CH)

72 Inventor: **O'Donnell, Margaret**
148 Allwood Road, Unit 4
Clifton, N.J. 07014(US)
Inventor: **Welton, Ann**
37 Maple Drive
North Caldwell, N.J. 07006(US)

74 Representative: **Kellenberger, Marcus, Dr. et**
al
Grenzacherstrasse 124 Postfach 3255
CH-4002 Basel(CH)

54 **Antiallergic combination.**

57 The invention relates to compositions comprising combinations of a PAF antagonist and a LTD₄ antagonist which combinations synergistically provide protection against allergic reactions such as antigen-induced death in mammals.

In another aspect, the invention relates to the use of the referred to combinations in the treatment of allergic reactions.

EP 0 469 477 A1

The present invention relates to compositions comprising the synergistic combination of platelet activating factor (PAF) antagonists with leukotriene D₄ (LTD₄) antagonists and the use thereof to treat allergic reactions.

The mechanism of pathogenesis associated with allergic reactions involves an antigen-induced release of a variety of allergic mediators from mast cells, basophils, and possibly macrophages. These mediators include histamine, leukotrienes and PAF all of which possess a broad spectrum of potent biological activities [Schleimer, R.P., Am. Rev. Respir. Dis., 133, 614-617 (1986) and Barnes, P.J., J. Allergy Clin. Immunol., 81, 152-160 (1988)]. Assessment of the precise contribution of each individual allergic mediator to the pathology of allergic reactions is unknown.

For example, allergic asthma involves the antigen-induced release of allergic mediators from mast cells which subsequently cause bronchoconstriction, mucus secretion, and changes in capillary permeability which, in combination with the production of chemotactic agents, leads to the influx into the lung of inflammatory cells. The mediators rapidly induce a narrowing and occluding of the bronchial lumen of the airways and, over time, the development of a chronic inflammatory condition in the lungs. Inflammation of the airways may be essential to the establishment of bronchial hyperresponsiveness which is the characteristic abnormality of asthma [Chung, K. F., Thorax, 41, 657-662 (1986)]. It has been difficult to identify the predominant mediator(s) of allergic asthma, but in recent years a number of lines of evidence strengthen the concept that the bronchoactive peptidoleukotrienes (LTC₄, LTD₄ and LTE₄) and PAF may be important in this disease process. LTD₄ and PAF have been identified in body fluids during an episode of asthma. They are both potent mediators of inflammation, and can reproduce some of the pathophysiological features of the disease when administered to animals and man [O'Donnell, M. in Therapeutic Approaches to Inflammatory Diseases, A. J. Lewis, N. S. Doherty and N. R. Ackerman, eds., Elsevier Science Publishing Co., New York, 169-193 (1989)].

Compounds which prevent the effects of the mediators are thus of interest in treating allergic reactions. However, since allergic reactions are complex processes that are probably not attributable to a single causative factor, a potent, specific, agent by itself may not be particularly effective in preventing allergic reactions.

Surprisingly, it has now been found that protection against allergic reactions such as antigen-induced death in mammals can be achieved by administering to such mammals an effective amount of a composition comprising a synergistic combination of a PAF antagonist and a LTD₄ antagonist.

In accordance with the invention, there can be used as the PAF antagonist component of a composition of the invention a known compound selected from the group consisting of:

5-{3-[4-(2-chlorophenyl)-9-methyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a]propynyl]phenanthridin-6(5H)-one; [1,4]diazepin-2-yl]-2-propynyl}phenanthridin-6(5H)-one;
 (R)-(2E,4E)-5-(4-methoxyphenyl)-N-[1-methyl-4-(3-pyridinyl)butyl]-2,4-decadienamide;
 3-[4-[(R)-2-[(methoxycarbonyl)oxy]-3-[(octadecylcarbamoyl)oxy]propoxy]butyl]thiazolium iodide;
 rac-4-{[6-(2-chlorophenyl)-8,9-dihydro-1-methyl-4H,7H-cyclopenta[4,5]thieno [3,2-f][1,2,4]triazolo[4,3-a][1,4]-diazepin-8-yl]carbonyl}morpholine;
 rac-N,N-dipropyl-6-[[6-(2-chlorophenyl)-8,9-dihydro-1-methyl-4H,7H-cyclopenta[4,5]thieno[3,2-f][1,2,4]-triazolo[4,3-a][1,4]diazepine-8-carboxamide;
 rac-6-(2-chlorophenyl)-8,9-dihydro-1-methyl-8-(4-morpholinyl)methyl-4H,7H-cyclopenta[4,5]thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine;
 4-(2-chlorophenyl)-9-methyl-2-{2-[4-(2-methylpropyl)phenyl]ethyl}-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]-diazepine;
 4-(2-chlorophenyl)-6,9-dimethyl-2-{2-[4-(2-methylpropyl)phenyl]ethyl}-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine;
 2,3-dihydro-5-{4-[2-(3,4,5-trimethoxyphenyl)ethyl]phenyl}imidazo[2,1-a]isoquinoline;
 (2S,5S)-2-[(3-methylsulfonyl-5-methoxy-4-propyloxy)phenyl]-5-(3,4,5-trimethoxyphenyl)tetrahydrofuran;
 (+)-2-(2-acetyl-6-methoxy-3,9-dioxo-4,8-dioxo-2,10-diazaoctacos-1-yl)-1-ethyl pyridinium chloride;
 (+)-N-(3-benzoylphenyl)-3-(3-pyridinyl)-1H,3H-pyrrolo[1,2-c]thiazole-7-carboxamide;
 1-O-hexadecyl-2(R,S)-O-ethyl-3-O-[7-(thiazolio-heptyl)]glycerol mesylate; and
 1-acetyl-4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)piperidine.

Preferred PAF antagonists comprise the following compounds:

5-{3-[4-(2-chlorophenyl)-9-methyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a]-[1,4]diazepin-2-yl]-2-propynyl}phenanthridin-6(5H)-one;
 (R)-(2E,4E)-5-(4-methoxyphenyl)-N-[1-methyl-4-(3-pyridinyl)butyl]-2,4-decadienamide;
 3-[4-[(R)-2-[(methoxycarbonyl)oxy]-3-[(octadecylcarbamoyl)oxy]propoxy]butyl]thiazolium iodide; and
 (2S,5S)-2-[(3-methylsulfonyl-5-methoxy-4-propyloxy)phenyl]-5-(3,4,5-trimethoxyphenyl)tetrahydrofuran.

Most preferred PAF antagonists are:

5-{3-[4-(2-chlorophenyl)-9-methyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a]-[1,4]diazepin-2-yl]-2-propynyl}phenanthridin-6(5H)-one; and
(R)-(2E,4E)-5-(4-methoxyphenyl)-N-[1-methyl-4-(3-pyridinyl)butyl]-2,4-decadienamide.

Also, in accordance with the invention, there can be used as the LTD₄ antagonist component of a composition a known compound selected from the group consisting of:

(E)-4-[3-[2-(4-cyclobutyl-2-thiazolyl)ethenyl]phenylamino]-2,2-diethyl-4-oxobutanoic acid;
rac-6-acetyl-7-[[5-(4-acetyl-3-hydroxy-2-propylphenoxy)pentyl]oxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid;

5-(3-(2-(7-chloroquinolin-2-yl)ethenyl)phenyl)-4,6-dithianonanedicarboxylic acid mono N,N-dimethylamide, sodium salt;

[3-[[2-methoxy-4-[[[(2-methylphenyl)sulfonyl]amino]carbonyl]phenyl]-methyl]-1-methyl-1H-indol-5-yl]-carbamic acid cyclopentyl ester;

(E)-4-[3-[2-(4-(1-methylethyl)-2-thiazolyl)ethenyl]phenylamino]-2,2-diethyl-4-oxobutanoic acid;

5-[[2-[[4-(2-quinolinylmethoxy)phenoxy]methyl]phenyl]methyl]-1H-tetrazole;

5-(7-(3-(quinolin-2-yl)methoxy)phenylmethoxy)-4-oxo-4H-1-benzopyran-2-yl)-1H-tetrazole;

3(S)-(2-(carboxethyl)thio)-3-(2-(8-phenyloctyl)phenyl)propanoic acid;

2(S)-hydroxy-3(R)-(2-(carboxethyl)thio)-3-(2-(8-phenyloctyl)phenyl)propanoic acid;

1-[2-hydroxy-3-propyl-4-[4-[2-[4-(1H-tetrazol-5-yl)butyl]-2H-tetrazol-5-yl]butoxy]phenyl]ethanone;

5-{3-[2(R)-(carboxyethylthio)-1(S)-hydroxypentadeca-3(E), 5(Z)-dienyl]phenyl}-1H-tetrazole

2-(3-(2-quinolinylmethoxy)phenylamino)benzoic acid;

1,1,1-trifluoro-N-[3-(2-quinolinylmethoxy)phenyl]methanesulfonamide;

N-[3-[3-(4-acetyl-3-hydroxy-2-propylphenoxy)-propoxy]-4-chloro-6-methylphenyl]-1H-tetrazole-5-carboxamide, sodium salt;

8-[4-(4-phenylbutoxy)benzoyl]amino-2-(tetrazol-5-yl)-4-oxo-4H-1-benzopyran;

N-[2,3-dihydro-3-(1H-tetrazol-5-yl)-1,4-benzodioxin-5-yl]-4-(4-phenylbutoxy)benzamide;

rac-7-[3-(4-acetyl-3-hydroxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid;

[[5-[[3-(4-acetyl-3-hydroxy-2-propylphenoxy)-propyl]thio]-1,3,4-thiadiazol-2-yl]thio]acetic acid;

9-(4-acetyl-3-hydroxy-2-propylphenoxy)methyl)-3-((1H)-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one;

6-(2-cyclohexylethyl)-[1,3,4]-thiadiazolo[3,2-a]-1,2,3-triazolo[4,5-d]pyrimidin-9(1H)-one; and

1-[[2-(1-methylethyl)pyrazolo[1,5-a]pyridin-3-yl]-2-methyl-1-propanone.

Preferred LTD₄ antagonists comprise the following compounds:

(E)-4-[3-[2-(4-cyclobutyl-2-thiazolyl)ethenyl]phenylamino]-2,2-diethyl-4-oxobutanoic acid;

rac-6-acetyl-7-[[5-(4-acetyl-3-hydroxy-2-propylphenoxy)pentyl]oxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid; and

5-(3-(2-(7-chloroquinolin-2-yl)ethenyl)phenyl)-4,6-dithianonanedicarboxylic acid mono N,N-dimethylamide, sodium salt.

The most preferred LTD₄ antagonist is:

(E)-4-[3-[2-(4-cyclobutyl-2-thiazolyl)ethenyl]phenylamino]-2,2-diethyl-4-oxobutanoic acid.

The synergistic compositions and methods of the invention due to their antiallergic activity, can be utilized for the treatment of allergic disorders such as seasonal rhinitis, perennial vasomotor rhinitis, acute urticaria, chronic urticaria, atopic dermatitis, contact dermatitis, pruritis, angioedema, conjunctivitis, chronic bronchitis, systemic anaphylaxis, serum sickness, bronchial asthma, food allergies, and related inflammatory diseases including inflammatory bowel disease, and psoriasis.

The anti-allergic effect of the compositions of the invention comprising combinations of PAF antagonists and LTD₄ antagonists may be demonstrated utilizing procedures described hereinafter.

A. Active Sensitization

Male (Hartley) guinea pigs (200-300 g) are sensitized with an intraperitoneal injection of ovalbumin (10 mg) in 1 ml of 0.9% w/v NaCl solution (saline) on days 1 and 3. The animals are used for the test on days 20-30 following the ovalbumin injections.

B. Pulmonary and Cardiovascular Mechanics

The sensitized guinea pigs are anesthetized with an intraperitoneal injection of urethane (2 g/kg), and polyethylene cannulae are introduced into both jugular veins and the right carotid artery. The arterial

catheter is connected to a pressure transducer for measurement to systemic blood pressure. The venous catheters are used for injection and infusion of the test composition.

The trachea is cannulated through an anterior cervical incision and positive pressure ventilation is applied with a Harvard pump set at a stroke volume of 4.5 ml and a rate of 40 breaths per minute. The respiratory muscles are paralyzed by periodic injections of succinylcholine chloride (1.2 mg/kg) given intravenously.

To measure transpulmonary pressure, a 12-gauge hypodermic needle attached via a short length of Tygon tubing to one side of a Statham differential pressure transducer is introduced through the fifth or sixth intercostal space into the intrapleural space near the sternum, leaving a pneumothorax. The other side of this transducer is connected to the side-arm of the tracheal cannula by a second Tygon tube of identical length and bore as the first. Airflow is measured with a Fleisch pneumotachograph head connected to a tracheal cannula and a second Statham differential pressure transducer. Electrical integration of the flow signal provides a recording of tidal volume.

Pulmonary resistance and dynamic lung compliance are calculated manually from the transpulmonary pressure, airflow, and volume signals as described by Amdur and Mead and the results are used to calibrate an on-line analog pulmonary computer [Am. J. Physiol., 192, 364-368 (1958)]. Blood pressure is recorded with a cannula inserted into the carotid artery. Heart rate is calculated by an on-line analog cardiovascular computer. All six parameters (airflow, transpulmonary pressure, tidal volume, resistance, compliance and blood pressure) are recorded simultaneously on a polygraph recorder and these parameters, plus heart rate and respiratory rate, are printed out on an electronic data terminal. After completion of surgery, a period of no less than 5 minutes is allowed for cardiopulmonary functions to stabilize. During this period, control values are recorded continuously.

C. Challenge

Following surgical preparation, the animals are pretreated with intravenous indomethacin (10 mg/kg) 22 minutes before ovalbumin challenge presumably to shunt arachidonic acid from the cyclooxygenase pathway into the lipoxygenase pathway for the formation of leukotrienes, and to eliminate the modulating effects of prostaglandins. Spontaneous breathing is arrested with succinylcholine (1.2 mg/kg, iv.) 7 minutes before antigen challenge. Treatment with succinylcholine results in a constant baseline ventilatory pressure of 6-8 cm H₂O. The bronchoconstrictor effects of histamine are abolished with pyrilamine (2.0 mg/kg, iv.), administered 6 minutes before antigen challenge. To potentiate the bronchoconstrictor effect of antigen, propranolol (0.1 mg/kg, i.v.) is administered before antigen challenge. [Anderson, W. H., et al., Br. J. Pharmacol., 78, 67-74 (1983)].

Various antagonists or control vehicle are administered intravenously through a cannula in the jugular vein 5 minutes before the animals are challenged with a maximum constrictory dose of ovalbumin (1 mg/kg) given intravenously.

The percent change in pulmonary resistance, dynamic lung compliance, and the systemic blood pressure and survival time are monitored for 120 minutes.

RESULTS

When the aforementioned model of anaphylaxis procedure was utilized to test various combinations of the invention, animals administered vehicle alone and challenged with antigen exhibited bronchoconstriction (approximately 550% increase in pulmonary resistance and 80% decrease in dynamic lung compliance), hypotension (55 mmHg decrease), and 83% mortality within 120 minutes. Table I contains the results obtained in this model with various combinations of the invention.

TABLE I

PROTECTION OF ANAPHYLACTIC DEATH IN SENSITIZED GUINEA
 5 FIG BY THE SYNERGISTIC COMPOSITIONS OF THE INVENTION

COMPOUNDS	DOSE mg/kg i.v.	COMPOUND	% SURVIVAL at 120 min.
ALONE *			
Control Vehicle			17
15 A	(1 mg/kg)	PAF antagonist	0
B	(1 mg/kg)	LTD ₄ antagonist	0
C	(1 mg/kg)	LTD ₄ antagonist	50
D	(10 mg/kg)	LTD ₄ antagonist	0
20 E	(1 mg/kg)	PAF antagonist	33
F	(1 mg/kg)	PAF antagonist	0
G	(1 mg/kg)	PAF antagonist	50
COMBINATIONS OF THE INVENTION *			
A+B			100
30 A+C			100
A+D			100
B+E			100
35 B+F			100
B+G			100
C+G			100

* COMPOUNDS

45 A = 5-[3-[4-(2-Chlorophenyl)-9-methyl-6H-thieno[3,2-f]1,2,4]triazolo-
 [4,3-a][1,4]diazepin-2-yl]-2-propynyl]phenanthridin-6(5H)-one.

50 B = (E)-4-[3-[2-(4-Cyclobutyl-2-thiazolyl)ethenyl]phenylamino]-2,2-diethyl-4-
 oxobutanoic acid.

C = 5-(3-(2-(7-Chloroquinolin-2-yl)ethenyl)phenyl)-4,6-dithianonane dicarboxylic acid, mono N,N-dimethylamide, sodium salt.

D = rac-6-Acetyl-7-[[5-(4-acetyl-3-hydroxy-2-propylphenoxy)pentyl]oxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid.

E = (R)-(2E,4E)-5-(4-Methoxyphenyl)-N-[1-methyl-4-(3-pyridinyl)butyl]-2,4-decadienamide.

F = 3-[4-[(R)-2-[(Methoxycarbonyl)oxy]-3-[(octadecylcarbamoyl)oxy]propoxy]-butyl]thiazolium iodide.

G = (2S,5S)-2-[(3-Methylsulfonyl-5-methoxy-4-propyloxy)phenyl]-5-(3,4,5-trimethoxyphenyl)tetrahydrofuran.

The ability of the LTD₄ antagonist, compound B [(E)-4-[3-[2(4-cyclobutyl-2-thiazolyl)ethenyl]-phenylamino]-2,2-diethyl-4-oxobutanoic acid] and the PAF antagonist, compound A [5-{3-[4-(2-chlorophenyl)-9-methyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-2-yl]-2-propynyl}phenanthridin-6-(5H)-one] to block antigen-induced bronchospasm, hypotension and death was evaluated at doses (1.0 mg/kg, i.v., 5 minute pretreatment) previously shown to completely inhibit bronchoconstriction induced by exogenous LTD₄ or PAF, respectively. Animals treated with compound B alone had a 0% survival rate at 120 minutes, whereas when the combination was given, there was 100% survival at 120 minutes.

The ability of the LTD₄ antagonist, compound C [5-(3-(2-(7-chloroquinolin-2-yl)ethenyl)phenyl)-4,6-dithianonedicarboxylic acid, mono N,N-dimethylamide, sodium salt] and the PAF antagonist, compound A to block antigen-induced bronchospasm, hypotension and death was evaluated at doses (1.0 mg/kg, i.v., 5 minute pretreatment) previously shown to completely inhibit bronchoconstriction induced by exogenous LTD₄ or PAF, respectively. Animals treated with compound C alone had a 50% survival rate at 120 minutes and those treated with compound A alone had a 0% survival rate at 120 minutes. However, when the combination was given, there was 100% survival at 120 minutes.

The ability of the LTD₄ antagonist, compound D [rac-6-acetyl-7-[[5-(4-acetyl-3-hydroxy-2-propylphenoxy)pentyl]oxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid] and the PAF antagonist, compound A to block antigen-induced bronchospasm, hypotension and death was evaluated at doses (10 mg/kg and 1.0 mg/kg, i.v., 5 minute pretreatment, respectively) previously shown to completely inhibit bronchoconstriction induced by exogenous LTD₄ or PAF, respectively. Animals treated with compound D or compound A alone had a 0% survival rate at 120 minutes, whereas when the combination was given, there was 100% survival at 120 minutes.

The ability of the LTD₄ antagonist, compound B and the PAF antagonist, compound E [(R) - (2E,4E) -5-(4-methoxyphenyl) -N- [1-methyl-4-(3-pyridinyl)butyl]-2,4-decadienamide] to block antigen-induced bronchospasm, hypotension and death was evaluated at doses (1.0 mg/kg, i.v., 5 minute pretreatment) previously shown to completely inhibit bronchoconstriction induced by exogenous LTD₄ or PAF, respectively. Animals treated with compound B alone had a 0% survival rate at 120 minutes and those treated with compound E alone had a 33% survival rate at 120 minutes. However, when the combination was given, there was 100% survival at 120 minutes.

The ability of the LTD₄ antagonist, compound B and the PAF antagonist, compound F [3-[4-[(R)-2-[(methoxycarbonyl)oxy]-3-[(octadecylcarbamoyl)oxy]propoxy]butyl]thiazolium iodide] to block antigen-induced bronchospasm, hypotension and death was evaluated at doses (1.0 mg/kg, i.v., 5 minute pretreatment) previously shown to completely inhibit bronchoconstriction induced by exogenous LTD₄ or PAF, respectively. Animals treated with compound B alone had a 0% survival rate at 120 minutes and those treated with compound F alone had a 0% survival rate at 120 minutes. However, when the combination was given, there was 100% survival at 120 minutes.

The ability of the LTD₄ antagonist, compound B and the PAF antagonist, compound G [(2S,5S)-2-[(3-methylsulfonyl-5-methoxy-4-propyloxy)phenyl]-5-(3,4,5-trimethoxyphenyl)tetrahydrofuran] to block antigen-

induced bronchospasm, hypotension and death was evaluated at doses (1.0 mg/kg, i.v., 5 minute pretreatment) previously shown to completely inhibit bronchoconstriction induced by exogenous LTD₄ or PAF, respectively. Animals treated with compound B alone had a 0% survival rate at 120 minutes and those treated with compound G alone had a 50% survival rate at 120 minutes. However, when the combination
5 was given, there was 100% survival at 120 minutes.

The ability of the LTD₄ antagonist, compound C and the PAF antagonist, compound G to block antigen-induced bronchospasm, hypotension and death was evaluated at doses (1.0 mg/kg, i.v., 5 minute pretreatment) previously shown to completely inhibit bronchoconstriction induced by exogenous LTD₄ or PAF, respectively. Animals treated with compound C alone had a 50% survival rate at 120 minutes and those
10 treated with compound G alone had a 50% survival rate at 120 minutes. However, when the combination was given, there was 100% survival at 120 minutes.

The compositions of the invention may be administered by any of the modes by which the individual components may be administered, for example, orally, intravenously or the like. The dosage regiment may be regulated according to the potency of individual compounds employed, the mode of administration, and
15 the needs of the host mammal depending on factors such as the degree and the severity of the disease state and age and general condition of the host mammal being treated.

The compositions of the invention can be administered in unit dosage forms such as tablets, capsules, pills, powders, granules, sterile parenteral solutions or suspensions, suppositories, transdermal compositions, aerosols and the like. Such dosage forms are prepared according to standard techniques in the art.

The dosages may be varied depending upon the requirements of the host, the severity of the condition being treated and the particular compound being employed in the composition. Determination of the proper dosage for a particular administration is within the skill of the art. Generally, treatment is initiated at lower dosages and increased as needed by small increments until the optimum effect is reached. Exemplary dosages are in the range of from 0.1 to 4.0 g per day of a combination of the invention. Exemplary dosages
20 of the PAF antagonist can be in the range of 0.01 to 2.0 g per day. Exemplary dosages of the LTD₄ antagonist can be in the range of 0.1 to 2.0 g per day. Generally, the PAF antagonist is present in a ratio of from about 0.01 to about 50 parts to one part of the LTD₄ antagonist, preferably in a ratio of from 0.1 to about 10 parts to one part of the LTD₄ antagonist. For convenience, the total daily dosage may be administered in divided doses.
25

30

EXAMPLARY FORMULATIONS COMPRISING THE INVENTION FOLLOW:

35

40

45

50

55

EXAMPLE 1TABLET FORMULATION

<u>Item</u>	<u>Ingredients</u>	<u>mg/Tablet</u>	
1	Compound A (1)	0.1 mg	250 mg
2	Compound B (1)	1.0 mg	250 mg
3	Croscarmellose Sodium*	6.4 mg	30 mg
4	Lactose	100 mg	255 mg
5	Avicel PH 101	20 mg	60 mg
6	Methocel E15**	1.5 mg	9 mg
7	Magnesium Stearate	<u>1.0 mg</u>	<u>6 mg</u>
		130 mg	860 mg

Method of Preparation:

1. Mix Items 1, 2, 3, 4, and 5 in a suitable mixer.
2. Granulate it with aqueous solution of Methocel E15.
3. Dry granulation at 45-50°C.
4. Pass the granulation through a suitable mill.
5. Add Item 7 and mix.
6. Compress the granulation on a suitable tablet press.

* Other disintegrants, such as starch derivatives, Cross-Povidone, Amberlite, or similar in action may be used alone or in combination to obtain desired disintegration characteristics.

** Binders, such as pregelatinized starch or polyvinyl-pyrrolidone, etc., may be used to obtain a desired binding. Wetting agents, such as Tweens, spans, or sodium lauryl sulfate may be added to obtain desired disintegration and dissolution characteristics.

(1) As used hereinafter-

Compound A is 5-{3-[4-(2-chlorophenyl)-9-methyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-2-yl]-2-propynyl} phenanthridin-6(5H)-one, and

Compound B is (E)-4-[3-[2-(4-cyclobutyl-2-thiazolyl)ethenyl]phenyl amino]-2,2-diethyl-4-oxobutanoic acid.

EXAMPLE 2CAPSULE FORMULATION

5	<u>Item</u>	<u>Ingredients</u>	<u>mg/Capsule</u>	
	1	Compound A	0.1 mg	250 mg
	2	Compound B	1.0 mg	250 mg
10	3	Pregelatinized Starch	5 mg	20 mg
	4	Modified Starch	5 mg	20 mg
	5	Talc	4.4 mg	20 mg
15	6	Lactose	84 mg	88 mg
	7	Magnesium Stearate	<u>0.5 mg</u>	<u>2 mg</u>
			100 mg	650 mg

20

Method of Preparation:

1. Mix Items 1, 2, 3, 4, and 6 and wet granulate with water.
2. Dry granulation at 45-50° C.
- 25
3. Mill through suitable screen using appropriate milling equipment.
4. Add Items 5 and 7 and mix for 5 minutes.
5. Fill into suitable capsules.

30

EXAMPLE 3SOFT GELATIN CAPSULE FORMULATION

35	<u>Item</u>	<u>Ingredients</u>	<u>mg/ml</u>	
	1	Compound A	0.1 mg	250 mg
	2	Compound B	1.0 mg	250 mg
40	3	Ethoxylated Fatty Acids	50 mg	250 mg
	4	Modified Starch	100 mg	100 mg
	5	Talc	1.0 ml	1.0 ml

45

Method of Preparation:

1. Add and mix Items 1 and 2 with Items 3 and 5.
2. Add Item 4 to the material from Step 1 and mix.
- 50
3. Add vegetable oil to the required amount.
4. Fill into a suitable capsule.

55

EXAMPLE 4ORAL SOLUTION FORMULATION

5

Item	Ingredients	<u>mg/5 ml</u>	
1	Compound A	0.1 mg	100 mg
2	Compound B	1.0 mg	100 mg
3	Alcohol, Anhydrous	10 mg	500 mg
4	PVP K-30	10 mg	500 mg
5	Aq. Sod. Hydroxide Solution	1.0 mg	100 mg
6	Sucrose	q.s.	q.s.
7	Flavoring Agent	q.s.	q.s.
8	PEG-400 q.s. to	5.0 ml	5.0 ml

20

Method of Preparation:

25

1. Dissolve Items 4,6, and 7 in alcohol and PEG-400.
2. Add Items 1 and 2 and mix thoroughly.
3. Add Item 5 to material from Step 2 and mix until Items 1 and 2 dissolve.
4. Add PEG-400 to the required volume.
5. Fill the solution into a suitable container.

30

35

40

45

50

55

EXAMPLE 5PARENTERAL FORMULATION

(for I. V. or IM use)

5

<u>Item</u>	<u>Ingredients</u>	<u>mg/ml</u>	
10	1 Compound A	0.1 mg	10 mg
	2 Compound B	1.0 mg	10 mg
	3 Benzyl Alcohol	10 mg	20 mg
15	4 Propylene Glycol*	0.4 ml	0.5 ml
	5 Emulphor EL620**	10 mg	50 mg
	6 Lactic Acid	0.01 ml	0.05 ml
20	7 Sod. Hydroxide pH to	6-7	6-7
	8 Water for Injection q.s. to	1.0 ml	1.0 ml

Method for Preparation:

25

1. Dissolve Items 3, 4, 5, and 6 in water for injection.
2. Add and dissolve Items 1 and 2 in the solution from Step 1.
3. Adjust pH if necessary.
4. Add water for injection to the required volume.

30

35

* Solvents or solubilizers, such as polyethylene glycols, alcohol, glycerine, Povidone, lecithin, sorbitan monooleate, or trioleate. Polysorbate 80 or 20 may be used in combination or alone to achieve the adequate solubility and stabilization.

40

** Powder for reconstitution may be formulated for appropriate solubilization or stabilization.

45

50

55

EXAMPLE 6INHALATION AEROSOL FORMULATION

(Suspension)

5

10

15

<u>Item</u>	<u>Ingredients</u>	<u>% w/w</u>
1	Compound A, Micronized	0.01
2	Compound B, Micronized	0.1
3	Sorbitan Trioleate	0.5
4	Freon 11	19.39
5	Freon 12	80
Total		100%

Method of Preparation:

25

1. Mix Items 1, 2, 3 and 4 and homogenize thoroughly.
2. Fill the concentrate suspension from Step 1 into a suitable can, place in valve, and crimp to seal container.
3. Pressure fill Item 5 to containers from Step 2.

EXAMPLE 7INHALATION AEROSOL FORMULATION

(Solution)

30

35

40

45

<u>Item</u>	<u>Ingredients</u>	<u>% w/w</u>
1	Compound A	0.01
2	Compound B	0.1
3	Ethyl Alcohol	10
4	Ascorbic Acid	0.5
5	Freon 114	20.0
6	Freon 12	68.5
Total		= 100%


Method of Preparation:

50

1. Dissolve Items 1, 2, and 4 in Item 3.
2. Fill solution from Step 1 into a suitable glass bottle, place in valve, and crimp to seal container.
3. Pressure fill Items 5 and 6 into the container from Step 2.

Claims

1. A pharmaceutical composition useful in the treatment of allergic disorders comprising an effective amount of a PAF antagonist selected from the group consisting of:



propynyl}phenanthridin-6(5H)-one;

(R)-(2E,4E)-5-(4-methoxyphenyl)-N-[1-methyl-4-(3-pyridinyl)butyl]-2,4-decadienamide;

3-[4-[(R)-2-[(methoxycarbonyl)oxy]-3-[(octadecylcarbamoyl)oxy] propoxy]butyl] thiazolium iodide;

rac-4-[[6-(2-chlorophenyl)-8,9-dihydro-1-methyl-4H,7H-cyclopenta [4,5]thieno[3,2-f][1,2,4]triazolo[4,3-a]-[1,4]diazepin-8-yl]carbonyl] morpholine;

rac-N,N-dipropyl-6-[[6-(2-chlorophenyl)-8,9-dihydro-1-methyl-4H,7H-cyclopenta[4,5]thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine-8-carboxamide;

rac-6-(2-chlorophenyl)-8,9-dihydro-1-methyl-8-(4-morpholinyl)methyl-4H,7H-cyclopenta[4,5]thieno[3,2-f]-[1,2,4]triazolo[4,3-a][1,4]diazepine;

4-(2-chlorophenyl)-9-methyl-2-{2-[4-(2-methylpropyl)phenyl]ethyl}-6H-thieno[3,2-f][1,2,4] triazolo[4,3-a]-[1,4]diazepine;

4-(2-chlorophenyl)-6,9-dimethyl-2-{2-[4-(2-methylpropyl)phenyl]ethyl}-6H-thieno[3,2-f][1,2,4] triazolo[4,3-a][1,4]diazepine;

2,3-dihydro-5-{4-[2-(3,4,5-trimethoxyphenyl)ethyl]phenyl}imidazo-[2,1-a]isoquinoline;

(2S,5S)-2-[[3-methylsulfonyl-5-methoxy-4-propyloxy]phenyl]-5-(3,4,5-trimethoxyphenyl)tetrahydrofuran;

(+)-2-(2-acetyl-6-methoxy-3,9-dioxo-4,8-dioxo-2,10-diazoctacos-1-yl)-1-ethyl pyridinium chloride;

(+)-N-(3-benzoylphenyl)-3-(3-pyridinyl)-1H,3H-pyrrolo[1,2-c]thiazole-7-carboxamide;

1-O-hexadecyl-2-(R,S)-O-ethyl-3-O-[7-(thiazolio-heptyl)]glycerol mesylate; and

1-acetyl-4-(8-chloro-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)piperidine,

and an effective amount of a LTD₄ antagonist selected from the group consisting of:

(E)-4-[3-[2-(4-cyclobutyl-2-thiazolyl)ethenyl]phenylamino]-2,2-diethyl-4-oxobutanoic acid;

rac-6-acetyl-7-[[5-(4-acetyl-3-hydroxy-2-propylphenoxy)pentyl]oxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid;

5-(3-(2-(7-chloroquinolin-2-yl)ethenyl)phenyl)-4,6-dithianonane dicarboxylic acid mono N,N-dimethylamide, sodium salt;

[3-[[2-methoxy-4-[[[(2-methylphenyl)sulfonyl]amino]carbonyl]phenyl] methyl]-1-methyl-1H-indol-5-yl]-carbamic acid cyclopentyl ester;

(E)-4-[3-[2-(4-(1-methylethyl)-2-thiazolyl)ethenyl]phenylamino]-2,2-diethyl-4-oxobutanoic acid;

5-[[2-[[4-(2-quinolinylmethoxy)phenoxy]methyl]phenyl]methyl]-1H-tetrazole;

5-(7-(3-(quinolin-2-yl)methoxy)phenylmethoxy)-4-oxo-4H-1-benzopyran-2-yl)-1H-tetrazole;

3 (S)-(2-(carboxethyl)thio-3-(2-(8-phenyloctyl)phenyl)propanoic acid;

2(S)-hydroxy-3(R)-(2-(carboxethyl)thio)-3-(2-(8-phenyloctyl)phenyl) propanoic acid;

1-[2-hydroxy-3-propyl-4-[4-[2-[4-(1H-tetrazol-5-yl)butyl]-2H-tetrazol-5-yl]butoxy] phenyl]ethanone;

5-{3-[2(R)-(carboxyethylthio)-1(S)-hydroxypentadeca-3(E),5(Z)-dienyl] phenyl}-1H-tetrazole;

2-(3-(2-quinolylmethoxy)phenylamino)benzoic acid;

1,1,1-trifluoro-N-[3-(2-quinolylmethoxy)phenyl]methanesulfonamide;

N-[3-[3-(4-acetyl-3-hydroxy-2-propylphenoxy)-propoxy]-4-chloro-6-methylphenyl]-1H-tetrazole-5-carboxamide, sodium salt;

8-[4-(4-phenylbutyloxy)benzoyl]amino-2-(tetrazol-5-yl)-4-oxo-4H-1-benzopyran;

N-[2,3-dihydro-3-(1H-tetrazol-5-yl)-1,4-benzodioxin-5-yl]-4-(4-phenyl butoxy) benzamide;

rac-7-[3-(4-acetyl-3-hydroxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid;

[[5-[[3-(4-acetyl-3-hydroxy-2-propylphenoxy)-propyl]thio]-1,3,4-thiadiazol-2-yl] thio]acetic acid;

(9-(4-acetyl-3-hydroxy-2-propylphenoxy)methyl)-3-((1H)-tetrazol-5-yl)-4H-pyrido [1,2-a]pyrimidin-4-one);

6-(2-cyclohexylethyl)-[1,3,4]-thiadiazolo[3,2-a]-1,2,3-triazolo[4,5-d] pyrimidin-9(1H)-one; and

1-[[2-(1-methylethyl)pyrazolo][1,5-a]pyridin-3-yl]-2-methyl-1-propanone,

and an inert pharmaceutical carrier.

2. A pharmaceutical composition, in accordance with claim 1, wherein the effective amount of the PAF antagonist is in the range of from about 0.01 to about 50 parts, to one part of the LTD₄ antagonist.

3. A pharmaceutical composition, in accordance with claim 1 or 2, wherein the PAF antagonist is selected from the group consisting of:

5-{3-[4-(2-chlorophenyl)-9-methyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4] diazepin-2-yl]-2-propynyl}phenanthridin-6(5H)-one;

(R)-(2E,4E)-5-(4-methoxyphenyl)-N-[1-methyl-4-(3-pyridinyl)butyl]-2,4-decadienamide;

3-[4-[(R)-2-[(methoxycarbonyl)oxy]-3-[(octadecylcarbamoyl)oxy] propoxy]butyl] thiazolium iodide; and

(2S,5S)-2-[(3-methylsulfonyl-5-methoxy-4-propyloxy)phenyl]-5-(3,4,5-trimethoxyphenyl)tetrahydrofuran,

and the LTD₄ antagonist is selected from the group consisting of:

(E)-4-[3-[2-(4-cyclobutyl-2-thiazolyl)ethenyl]phenylamino]-2,2-diethyl-4-oxobutanoic acid;

rac-6-acetyl-7-[[5-(4-acetyl-3-hydroxy-2-propylphenoxy)pentyl]oxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid; and

5-(3-(2-(7-chloroquinolin-2-yl)ethenyl)phenyl)-4,6-dithianonane dicarboxylic acid mono N,N-dimethylamide, sodium salt.

4. A pharmaceutical composition, in accordance with claim 3, wherein the PAF antagonist is selected from the group consisting of:

5-{3-[4-(2-chlorophenyl)-9-methyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a] [1,4]diazepin-2-yl]-2-propynyl}phenanthridin-6(5H)-one; and

(R)-(2E,4E)-5-(4-methoxyphenyl)-N-[1-methyl-4-(3-pyridinyl)butyl]-2,4-decadienamide,

and the LTD₄ antagonist is

(E)-4-[3-[2-(4-cyclobutyl-2-thiazolyl)ethenyl]phenylamino]-2,2-diethyl-4-oxobutanoic acid.

5

5. A pharmaceutical composition, in accordance with claim 4, wherein the PAF antagonist is

5-{3-[4-(2-chlorophenyl)-9-methyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a] [1,4]diazepin-2-yl]-2-propynyl}phenanthridin-6(5H)-one,

10

and the LTD₄ antagonist is

(E)-4-[3-[2-(4-cyclobutyl-2-thiazolyl)ethenyl]phenylamino]-2,2-diethyl-4-oxobutanoic acid.

15

6. A pharmaceutical composition, in accordance with claim 4, wherein the PAF antagonist is

(R)-(2E,4E)-5-(4-methoxyphenyl)-N-[1-methyl-4-(3-pyridinyl)butyl]-2,4-decadienamide,

and the LTD₄ antagonist is

20

(E)-4-[3-[2-(4-cyclobutyl-2-thiazolyl)ethenyl]phenylamino]-2,2-diethyl-4-oxobutanoic acid.

7. The use of a composition in accordance with any one of claims 1 to 6 in the control or prevention of allergic disorders or for the manufacture of medicaments against allergic disorders.

25

30

35

40

45

50

55



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 91 11 2577

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	CHEMICAL ABSTRACTS, vol. 110, no. 21, 22nd May 1989, page 586, abstract no. 190981f, Columbus, Ohio, US; R. PATTERSON et al.: "Effects of combined receptor antagonists of leukotriene D ₄ (LTD ₄) and platelet-activating factor (PAF) on rhesus airway responses to LTD ₄ , PAF and antigen", & INT. ARCH. ALLERGY APPL. IMMUNOL. 1989, 88(4), 462-70 * Abstract *	1-7	A 61 K 31/55 A 61 K 31/44 // (A 61 K 31/55 A 61 K 31:425) (A 61 K 31/44 A 61 K 31:425)
A	EP-A-0 345 931 (SCHERING CORP.) -----	1-7	
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			A 61 K
The present search report has been drawn up for all claims			
Place of search		Date of completion of search	Examiner
The Hague		04 November 91	BRINKMANN C.
CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document			